



GROW SCHEDULE



	VEGETATIVE PHASE				GENERATIVE PHASE						
	ROOTING		1		I I			П	III	IV	
DAYS AND WEEKS	D1-5	D6-25	W1	W2	W3	W4	W5	W6	W7	W8	W9
LIGHT HOURS	18	18	12	12	12	12	12	12	12	12	12
	Dosages in ml per 10 L										
CANNA COCO A	26 - 33	26 - 33	29 - 37	29 - 37	29 - 37	29 - 37	29 - 37	29 - 37	26 - 33	26 - 33	
CANNA COCO B	26 - 33	26 - 33	29 - 37	29 - 37	29 - 37	29 - 37	29 - 37	29 - 37	26 - 33	26 - 33	
CANNA RHIZOTONIC XP	40	20	20	20	5	5	5	5	5		
CANNAZYM		25	25	25	25	25	25	25	25	25 - 50 *	25 - 50
CANNABOOST			20	20	20 - 40 **	20 - 40	20 - 40	20 - 40	20 - 40	20 - 40	20 - 40
CANNA PK 13/14								15			
	m\$/cm										
EC WATER	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
EC NUTRITION	1.4 - 1.8	1.4 - 1.8	1.6 - 2.0	1.6 - 2.0	1.6 - 2.0	1.6 - 2.0	1.6 - 2.0	2.0 - 2.5	1.4 - 1.8	1.4 - 1.8	
EC TOTAL	1.8 - 2.2	1.8 - 2.2	2.0 - 2.4	2.0 - 2.4	2.0 - 2.4	2.0 - 2.4	2.0 - 2.4	2.4 - 2.9	1.8 - 2.2	1.8 - 2.2	0.4

This cultivation schedule is intended for plants that develop flowers or fruits, starting from a one-week-old cutting or seedling. The duration of each phase may vary depending on the species.

- * Double CANNAZYM dosage to 50 ml/10 L, if substrate is reused.
- ** Increase CANNABOOST to a maximum of 40 ml/10 L through flowering for extra strength.

For a longer growth phase, repeat the recommended dosages from Rooting period II (D6-25). For a longer flowering phase, repeat the recommended dosages from generative Phase I (W3-W5).

Add CANNA PK 13/14 in the period 3-4 weeks before harvest.



