



## GROW SCHEDULE



	VEGETATIVE PHASE				GENERATIVE PHASE						
	ROO	ROOTING		l l		l l			III	IV	
DAYS AND WEEKS	D1-5	D6-25	W1	W2	W3	W4	W5	W6	W7	W8	W9
LIGHT HOURS	18	18	12	12	12	12	12	12	12	12	12
	Dosages in ml per 10 L										
CANNA SUBSTRA VEGA A (HW)	24 - 31	28 - 35	28 - 35	28 - 35							
CANNA SUBSTRA VEGA B (HW)	24 - 31	28 - 35	28 - 35	28 - 35							
CANNA SUBSTRA FLORES A (HW)					28 - 35	28 - 35	28 - 35	28 - 35	24 - 32	21 - 28	
CANNA SUBSTRA FLORES B (HW)					28 - 35	28 - 35	28 - 35	28 - 35	24 - 32	21 - 28	
CANNA RHIZOTONIC XP	40	20	20	20	5	5	5	5	5		
CANNAZYM		25	25	25	25	25	25	25	25	25 - 50 <sup>*</sup>	25 - 50
CANNABOOST			20	20	20 - 40 **	20 - 40	20 - 40	20 - 40	20 - 40	20 - 40	20 - 40
CANNA PK 13/14								15			
	m\$/cm										
EC WATER	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
EC NUTRITION	1.5 - 1.9	1.7 - 2.1	1.7 - 2.1	1.7 - 2.1	1.7 - 2.1	1.7 - 2.1	1.7 - 2.1	2.0 - 2.4	1.4 - 1.9	1.2 - 1.7	
EC TOTAL	2.1 - 2.5	2.3 - 2.7	2.3 - 2.7	2.3 - 2.7	2.3 - 2.7	2.3 - 2.7	2.3 - 2.7	2.6 - 3.0	2.0 - 2.5	1.8 - 2.3	0.6

This cultivation schedule is intended for plants that develop flowers or fruits, starting from a one-week-old cutting or seedling. The duration of each phase may vary depending on the species.

- \* Double CANNAZYM dosage to 50 ml/10 L, if substrate is reused.
- \*\* Increase CANNABOOST to a maximum of 40 ml/10 L through flowering for extra strength.

For a longer growth phase, repeat the recommended dosages from Rooting period II (D6-25). For a longer flowering phase, repeat the recommended dosages from generative Phase I (W3-W5).

Add CANNA PK 13/14 in the period 3-4 weeks before harvest.



